



SHANDONG QILU KING-PHAR PHARMACEUTICAL CO.,LTD.

山东齐发药业有限公司

ADD:243 GONGYE NORTH ROAD, JINAN, 250100, CHINA

TEL: +86 531 83127765

FAX: +86 531 83127769

E-mail: qilupy@qilu-pharma.com <http://www.qilupharma.com>

Monensin Premix

Brief Introduction: Monensin anticoccidiostat with wide spectrum belongs to polyether antibiotic, and is one of most typical veterinary of anticoccidiostat that is first used and most marketable in the world. Gram-positive and fungus on vegetable are sensitive to it. It can inhibits coccid's growth and finally kill it, especially to eimeria tenella coccidial.

Mechanism:

1.The special structure of monensin furnishes its character of high affinity with earth metal iron, such as K, Na iron etc, which enhances the permeant ability of necessary cation passing through the ether barrier in the membrane, resulting in the iron balance changing in coccid and the death of it, so monensin is a typical ionophore antibiotic.

2. Monensin can improve the utility of crude protein, crude fatty acid, crud cellulose in feedstuff, so it can improve the conversion ratio of feedstuff.

3. Effectively improving the content of propionic acid in the stomach of ruminant and reducing the generation of acetic, which comes to a much more conversion of feedstuff.

Specification: 20%, 40% or other content required by the clients.

Indication: First, anticoccidiostat, it can reduce incidence of a disease and bring down mortality, improve the conversion ratio of feedstuff and promote the growth of poultry and livestock. Second, it can enhance the content of propionic acid, promote feedstuff to convert largely.

Administration and dosage: First dilute monensin with proper amount of feedstock, until it is mixed homogeneously, then further dilution and mixing until mixture containing the required content:

1. For treating coccidiosis in poultry and other animals:90-110ppm(this is 90g-110g monensin per ton of feedstock.)
2. Cattle breeding:30ppm(30g monensin per ton of feedstock.) or feeding as per 100mg monensin per cattle per day.

Caution:

1. It is prohibited to feed chickens during the term of laying eggs.
2. It is toxic to turkey and horse.
3. It is prohibited to use with tilamuin or other anticoccidiostat together.
4. Halting period: 3 days.
5. When blending, avoid contacting with skins and eyes.

Storage Avoid sunlight, store in dry place.

Term of validity Two years

Package 25kg per sack.

Exclusive Distributor :

JNS **JNS TECHNOLOGY**
.....Juvenile Nutritional Support

Uttaran, House # 11 (2nd Floor),
Road # 14, Sector # 01, Uttara,
Dhaka-1230, Bangladesh
Tel : +88 02 7911083, Cell : +88 01711447017
Email : jakirbncc@gmail.com